



MARKSCHEME

November 2014

**MATHEMATICS
CALCULUS**

Higher Level

Paper 3

*This markscheme is **confidential** and for the exclusive use of examiners in this examination session.*

*It is the property of the International Baccalaureate and must **not** be reproduced or distributed to any other person without the authorization of the IB Assessment Centre.*

Instructions to Examiners

Abbreviations

- M** Marks awarded for attempting to use a correct **Method**; working must be seen.
- (M)** Marks awarded for **Method**; may be implied by **correct** subsequent working.
- A** Marks awarded for an **Answer** or for **Accuracy**; often dependent on preceding **M** marks.
- (A)** Marks awarded for an **Answer** or for **Accuracy**; may be implied by **correct** subsequent working.
- R** Marks awarded for clear **Reasoning**.
- N** Marks awarded for **correct** answers if **no** working shown.
- AG** Answer given in the question and so no marks are awarded.

Using the markscheme

1 General

Mark according to RM™ Assessor instructions and the document “**Mathematics HL: Guidance for e-marking November 2014**”. It is essential that you read this document before you start marking. In particular, please note the following:

- Marks must be recorded using the annotation stamps. Please check that you are entering marks for the right question.
- If a part is **completely correct**, (and gains all the “must be seen” marks), use the ticks with numbers to stamp full marks.
- If a part is completely wrong, stamp **A0** by the final answer.
- If a part gains anything else, it **must** be recorded using **all** the annotations.
- All the marks will be added and recorded by RM™ Assessor.

2 Method and Answer/Accuracy marks

- Do **not** automatically award full marks for a correct answer; all working **must** be checked, and marks awarded according to the markscheme.
- It is not possible to award **M0** followed by **A1**, as **A** mark(s) depend on the preceding **M** mark(s), if any.
- Where **M** and **A** marks are noted on the same line, eg **M1A1**, this usually means **M1** for an **attempt** to use an appropriate method (eg substitution into a formula) and **A1** for using the **correct** values.
- Where the markscheme specifies **(M2)**, **N3**, etc., do **not** split the marks.
- Once a correct answer to a question or part-question is seen, ignore further working.

3 N marks

*Award **N** marks for **correct** answers where there is **no** working.*

- Do **not** award a mixture of **N** and other marks.
- There may be fewer **N** marks available than the total of **M**, **A** and **R** marks; this is deliberate as it penalizes candidates for not following the instruction to show their working.

4 Implied marks

*Implied marks appear in **brackets eg (M1)**, and can only be awarded if **correct** work is seen or if implied in subsequent working.*

- Normally the correct work is seen or implied in the next line.
- Marks **without** brackets can only be awarded for work that is **seen**.

5 Follow through marks

*Follow through (**FT**) marks are awarded where an incorrect answer from one **part** of a question is used correctly in **subsequent** part(s). To award **FT** marks, **there must be working present** and not just a final answer based on an incorrect answer to a previous part.*

- If the question becomes much simpler because of an error then use discretion to award fewer **FT** marks.
- If the error leads to an inappropriate value (eg $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).
- Within a question part, once an error is made, no further **dependent A** marks can be awarded, but **M** marks may be awarded if appropriate.
- Exceptions to this rule will be explicitly noted on the markscheme.

6 Mis-read

*If a candidate incorrectly copies information from the question, this is a mis-read (**MR**). A candidate should be penalized only once for a particular mis-read. Use the **MR** stamp to indicate that this has been a misread. Then deduct the first of the marks to be awarded, even if this is an **M** mark, but award all others so that the candidate only loses one mark.*

- If the question becomes much simpler because of the **MR**, then use discretion to award fewer marks.
- If the **MR** leads to an inappropriate value (eg $\sin \theta = 1.5$), do not award the mark(s) for the final answer(s).

7 Discretionary marks (*d*)

*An examiner uses discretion to award a mark on the rare occasions when the markscheme does not cover the work seen. In such cases the annotation **DM** should be used and a brief **note** written next to the mark explaining this decision.*

8 Alternative methods

Candidates will sometimes use methods other than those in the markscheme. Unless the question specifies a method, other correct methods should be marked in line with the markscheme. If in doubt, contact your team leader for advice.

- Alternative methods for complete questions are indicated by **METHOD 1, METHOD 2, etc.**
- Alternative solutions for part-questions are indicated by **EITHER . . . OR.**
- Where possible, alignment will also be used to assist examiners in identifying where these alternatives start and finish.

9 Alternative forms

Unless the question specifies otherwise, **accept** equivalent forms.

- As this is an international examination, accept all alternative forms of **notation**.
- In the markscheme, equivalent **numerical** and **algebraic** forms will generally be written in brackets immediately following the answer.
- In the markscheme, **simplified** answers, (which candidates often do not write in examinations), will generally appear in brackets. Marks should be awarded for either the form preceding the bracket or the form in brackets (if it is seen).

Example: for differentiating $f(x) = 2\sin(5x - 3)$, the markscheme gives:

$$f'(x) = (2\cos(5x - 3))5 \quad (=10\cos(5x - 3)) \quad \text{AI}$$

Award **AI** for $(2\cos(5x - 3))5$, even if $10\cos(5x - 3)$ is not seen.

10 Accuracy of Answers

Candidates should **NO LONGER** be penalized for an accuracy error (**AP**).

If the level of accuracy is specified in the question, a mark will be allocated for giving the answer to the required accuracy. When this is not specified in the question, all numerical answers should be given exactly or correct to three significant figures. Please check work carefully for **FT**.

11 Crossed out work

If a candidate has drawn a line through work on their examination script, or in some other way crossed out their work, do not award any marks for that work.

12 Calculators

A GDC is required for paper 3, but calculators with symbolic manipulation features (for example, TI-89) are not allowed.

Calculator notation

The Mathematics HL guide says:

Students must always use correct mathematical notation, not calculator notation.

Do **not** accept final answers written using calculator notation. However, do not penalize the use of calculator notation in the working.

13 More than one solution

Where a candidate offers two or more different answers to the same question, an examiner should only mark the first response unless the candidate indicates otherwise.

1. (a) $\int_1^{\infty} x^{-0.5} dx$ *M1*

$$= \lim_{H \rightarrow \infty} \left[2x^{0.5} \right]_1^H$$
 A1

Note: Accept $\left[2x^{0.5} \right]_1^{\infty}$.

this is not finite so series is divergent *R1*

[3 marks]

Note: Accept equivalent eg $\rightarrow \infty$, or “limit does not exist”.

If lower limit is not equal to 1 award *M0A0*, but the *R1* can still be awarded if the final reasoning is correct.

(b) (i) applying the ratio test *M1*

$$\lim_{n \rightarrow \infty} \left| \frac{(x+1)^{n+1}}{2^{n+1}(n+1)^{0.5}} \times \frac{2^n n^{0.5}}{(x+1)^n} \right|$$
 A1

$$\lim_{n \rightarrow \infty} \left| \frac{(x+1)n^{0.5}}{2(n+1)^{0.5}} \right| = \left| \frac{(x+1)}{2} \right|$$
 A1

Note: Do not penalize the absence of limits and modulus signs.

converges if $\left| \frac{x+1}{2} \right| < 1 \Rightarrow -1 < \frac{(x+1)}{2} < 1$ *M1*

$\Rightarrow -3 < x < 1$ *A1*

Note: Accept $-2 < x+1 < 2$.

(ii) considering end points *M1*

when $x = -3$, series is $\sum_{n=1}^{\infty} \frac{(-1)^n}{n^{0.5}}$ *A1*

$\frac{1}{n^{0.5}}$ is a decreasing sequence with limit zero, *R1*

so series converges by alternating series test *R1*

when $x = 1$, series is $\sum_{n=1}^{\infty} \frac{1}{n^{0.5}}$ which diverges by part (a) or

p -series *A1*

Note: This *A1* is for both the reasoning and the statement it diverges.

interval of convergence is $-3 \leq x < 1$ *A1*

[11 marks]

Total [14 marks]

2. (a) integrating factor $e^{\int \frac{1}{t} dt} = e^{-\ln t} \left(= \frac{1}{t} \right)$ **M1A1**

$$\frac{x}{t} = \int -\frac{2}{t^2} dt = \frac{2}{t} + c$$
A1A1

Note: Award **A1** for $\frac{x}{t}$ and **A1** for $\frac{2}{t} + c$.

$$x = 2 + ct$$
AG
[4 marks]

(b) given continuity at $x = 5$

$$5c + 2 = 16 - \frac{35}{5} \Rightarrow c = \frac{7}{5}$$
M1A1
[2 marks]

(c) (i) 2 **A1**

(ii) any value ≥ 16 **A1**

Note: Accept values less than 16 if fully justified by reference to the maximum age for a dog.

[2 marks]

continued ...

Question 2 continued

$$(d) \lim_{h \rightarrow 0^-} \left(\frac{\frac{7}{5}(5+h) + 2 - \frac{7}{5}(5) - 2}{h} \right) = \frac{7}{5} \quad \text{M1A1}$$

$$\lim_{h \rightarrow 0^+} \left(\frac{16 - \frac{35}{5+h} - 16 + \frac{35}{5}}{h} \right) \left(= \lim_{h \rightarrow 0^+} \left(\frac{-35}{5+h} + 7 \right) \right) \quad \text{M1}$$

$$= \lim_{h \rightarrow 0^+} \left(\frac{\frac{-35+35+7h}{(5+h)}}{h} \right) = \lim_{h \rightarrow 0^+} \left(\frac{7}{5+h} \right) = \frac{7}{5} \quad \text{M1A1}$$

both limits equal so differentiable at $t = 5$ RIAG

Note: The limits $t \rightarrow 5$ could also be used.

For each value of $\frac{7}{5}$ obtained by standard differentiation award **A1**.

To gain the other 4 marks a rigorous explanation must be given on how you can get from the left and right hand derivatives to the derivative.

Note: If the candidate works with t and then substitutes $t = 5$ at the end award as follows


First **M1** for using formula with t in the linear case, **A1** for $\frac{7}{5}$

Award next 2 method marks even if $t = 5$ not substituted, **A1** for $\frac{7}{5}$

[6 marks]

Total [14 marks]

3. (a) and (b)



- (a) *AI* for 4 parallel straight lines with a positive gradient
AI for correct y intercepts

AI

AI

[2 marks]

- (b) *AI* for passing through $(0, 1)$ with positive gradient less than 2
AI for stationary point on $y = 2x$
AI for negative gradient on both of the other 2 isoclines

*AI**AI**AI*

[3 marks]

- (c) the isocline is perpendicular to C

R1

[1 mark]

- (d) $y_{n+1} = y_n + 0.1(y_n - 2x_n) (= 1.1y_n - 0.2x_n)$

(*M1*)(*AI*)

Note: Also award *M1AI* if no formula seen but y_2 is correct.

$$y_0 = 1, y_1 = 1.1, y_2 = 1.19, y_3 = 1.269, y_4 = 1.3359$$

(*M1*)

$$y_5 = 1.39 \text{ to 3sf}$$

AI

Note: *M1* is for repeated use of their formula, with steps of 0.1.

Note: Accept 1.39 or 1.4 only.

[4 marks]

Total [10 marks]

4. (a) $r = -x^2, \quad S = \frac{1}{1+x^2}$

AIAG

[1 mark]

(b) $\frac{1}{1+x^2} = 1 - x^2 + x^4 - x^6 + \dots$

EITHER

$$\int \frac{1}{1+x^2} dx = \int 1 - x^2 + x^4 - x^6 + \dots dx$$

M1

$$\arctan x = c + x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$$

AI

Note: Do not penalize the absence of c at this stage.

when $x = 0$ we have $\arctan 0 = c$ hence $c = 0$

M1A1

OR

$$\int_0^x \frac{1}{1+t^2} dt = \int_0^x 1 - t^2 + t^4 - t^6 + \dots dt$$

M1A1A1

Note: Allow x as the variable as well as the limit.
M1 for knowing to integrate, **AI** for each of the limits.

$$[\arctan t]_0^x = \left[t - \frac{t^3}{3} + \frac{t^5}{5} - \frac{t^7}{7} + \dots \right]_0^x$$

AI

$$\text{hence } \arctan x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$$

AG

[4 marks]

(c) applying the MVT to the function f on the interval $[x, y]$

M1

$$\frac{f(y) - f(x)}{y - x} = f'(c) \quad (\text{for some } c \in]x, y[)$$

AI

$$\frac{f(y) - f(x)}{y - x} > 0 \quad (\text{as } f'(c) > 0)$$

R1

$$f(y) - f(x) > 0 \quad \text{as } y > x$$

R1

$$\Rightarrow f(y) > f(x)$$

AG

[4 marks]

Note: If they use x rather than c they should be awarded **M1A0R0**, but could get the next **R1**.

continued...

Question 4 continued

(d) (i) $g(x) = x - \arctan x \Rightarrow g'(x) = 1 - \frac{1}{1+x^2}$ **A1**

this is greater than zero because $\frac{1}{1+x^2} < 1$ **R1**

so $g'(x) > 0$ **AG**

(ii) (g is a continuous function defined on $[0, b]$ and differentiable on $]0, b[$ with $g'(x) > 0$ on $]0, b[$ for all $b \in \mathbb{R}$)

(If $x \in [0, b]$ then) from part (c) $g(x) > g(0)$ **M1**

$x - \arctan x > 0 \Rightarrow \arctan x < x$ **M1**

(as b can take any positive value it is true for all $x > 0$) **AG**

[4 marks]

(e) let $h(x) = \arctan x - \left(x - \frac{x^3}{3}\right)$ **M1**

(h is a continuous function defined on $[0, b]$ and differentiable on $]0, b[$ with $h'(x) > 0$ on $]0, b[$)

$h'(x) = \frac{1}{1+x^2} - (1 - x^2)$ **A1**

$= \frac{1 - (1 - x^2)(1 + x^2)}{1 + x^2} = \frac{x^4}{1 + x^2}$ **M1A1**

$h'(x) > 0$ hence (for $x \in [0, b]$) $h(x) > h(0) (= 0)$ **R1**

$\Rightarrow \arctan x > x - \frac{x^3}{3}$ **AG**

[5 marks]

Note: Allow correct working with $h(x) = x - \frac{x^3}{3} - \arctan x$.

continued ...

Question 4 continued

(f) use of $x - \frac{x^3}{3} < \arctan x < x$

MI

choice of $x = \frac{1}{\sqrt{3}}$

AI

$$\frac{1}{\sqrt{3}} - \frac{1}{9\sqrt{3}} < \frac{\pi}{6} < \frac{1}{\sqrt{3}}$$

MI

$$\frac{8}{9\sqrt{3}} < \frac{\pi}{6} < \frac{1}{\sqrt{3}}$$

AI

Note: Award final ***AI*** for a correct inequality with a single fraction on each side that leads to the final answer.

$$\frac{16}{3\sqrt{3}} < \pi < \frac{6}{\sqrt{3}}$$

AG

[4 marks]

Total [22 marks]